

MOSAIC

Deliverable Report

D.2.1.1

Document identifier	MOSAIC_D2.1.1
Due Date	31 January 2026
Document Date	14 January 2026
Deliverable Title	Field and lab experiments database - Field and lab data published and documented in open-access data repositories
Level of Dissemination	Public
Work Package	2
Lead Partner	PP3 UNITO
Author(s)	Nicolò Anselmetto, Raffaella Marzano, Matteo Garbarino

Version	Date	Author(s)	Changes
2	14 January 2026	N. Anselmetto, R. Marzano, M. Garbarino	V2

This project is co-funded by the European Union through the Interreg Alpine Space programme.

Alpine Space

MOSAIC

Table of Contents

Table of Contents	1
Executive Summary	4
1. Introduction.....	5
1.1 Purpose of Deliverable	5
1.2 Deliverable Overview	6
1.3 Data Management, FAIR Principles, and Accessibility	7
1.4 Structure of the Deliverable.....	9
1.5 Related Documents	9
2. Field and Lab Experiments Database	10
2.1 Chapter introduction	10
2.2 Structure of the database table.....	10
2.3 The Database	12
3. Standardized Metadata Submission Framework.....	23
3.1 Introduction	23
3.2 Submission Form Fields	24
3.2.1 Identification	24
3.2.2 Description	25
3.2.3 Spatial and temporal scales	26

Alpine Space

MOSAIC

3.2.4 Methods	27
3.2.5 References.....	28
4. Conclusions and Outlook.....	29
5. References.....	32

Alpine Space

MOSAIC

Executive Summary

The deliverable **D.2.1.1 “Field and lab experiments database”** encompasses field data, laboratory data, derived products, and models developed or used within the project and published and documented in open-access data repositories. The database is designed around the FAIR principles (Findability, Accessibility, Interoperability, and Reusability) to support data sharing, transparency, and reuse, and consists of publicly accessible data through several online repositories, such as Figshare, Dryad, or Zenodo. Notably, several data curated within the project will be made publicly accessible through the MOSAIC WebAtlas, accessible at <https://alpineresilience.org/>.

The deliverable report is structured in five chapters. The first chapter *Introduction* provides an overview of the deliverable, its purpose, and related documents. The second chapter, *Field and Lab Experiments Database*, is the basic table with data records and models currently included in the database, describing their thematic scope, data type, responsible partners, and access links. The third chapter, *Standardized Metadata Submission Framework*, introduces the standardized submission form and metadata framework to guide partners in contributing additional datasets and models in a consistent, interoperable, and FAIR-compliant manner. The fourth and fifth chapters encompass conclusions and references, respectively.

Alpine Space

MOSAIC

1. Introduction

1.1 Purpose of Deliverable

This deliverable documents the release of the MOSAIC database developed under **Work Package 2 (WP2): NAZCA – *Natural haZards modelling platform for analysing climate change (CC) compound events on Alpine Space (AS) protective forests***. The overarching goal of WP2 is to provide a modelling platform and associated datasets integrating natural hazard (e.g. rockfalls, avalanches, wildfires, windthrows) and risk models upgraded to account for the consequences of CC - such as changes in species suitability - and the effects of compound events on forest composition, structure, and protective functions of the AS.

Within this framework, **Activity 2.1** focuses on generating missing empirical evidence through **field surveys and field and laboratory experiments** addressing key processes related to CC-driven compound events threatening AS forests. These include, among others, mechanical properties of trees, fire and windthrow susceptibility, snowpack characteristics, regeneration dynamics, and the prioritisation of post-disturbance management practices. The data produced enable an improved understanding of CC impacts on hazard behaviour, forest and disturbance dynamics, and ultimately on the effectiveness of protective forests as **Ecosystem-based solutions for Disaster Risk Reduction (Eco-DRR)**.

The objective of this deliverable is to organise, document, and publish these datasets and associated models in a structured, open-access database designed around the **FAIR principles** (Wilkinson et al., 2016). By ensuring standardized documentation, persistent identification, and open accessibility, the deliverable supports the long-term availability, traceability, and usability of project outputs for scientific research, technical applications, and policy-oriented decision-

MOSAIC

making, and provides a foundation for their integration into the MOSAIC WebAtlas and modelling platform.

1.2 Deliverable Overview

Climate change (CC) is increasingly responsible for the rise in climate-related disasters across Europe, resulting in substantial socio-economic losses and impacts on human well-being (IPCC 2022). Many of these disasters are driven by compound and cascading events, i.e. the interaction of multiple climate-related hazards occurring simultaneously or sequentially, which together amplify socio-ecological risks (Cutter 2018). Addressing such risks requires **integrated, spatially explicit data and models at different scales** that capture interactions between CC, natural hazards, forest ecosystems, and human systems.

In this context, forests play a dual role. They provide essential ecosystem services that support climate change mitigation and adaptation, and in the AS they represent a key **Eco-DRR** by protecting people and infrastructure against natural hazards (Dorren et al. 2004, Moos et al. 2023). At the same time, forest health, structure, and protective functions are themselves increasingly threatened by CC-driven disturbances such as wildfires, windstorms, pest outbreaks, and changes in snow and hydrological regimes (Bastit 2023). Ensuring hazard-resilient and sustainable forest management therefore requires proactive, transnational strategies supported by harmonised data and models.

This deliverable presents a **harmonised inventory of datasets, models, and derived products** generated within MOSAIC or sourced from relevant external initiatives. These resources directly support the development and implementation of NAZCA, which integrates climate-related compound events into natural hazard and risk models adapted to Alpine-specific conditions (e.g. topography, forest types, and management systems). The database contributes to assessing

Alpine Space

MOSAIC

socio-ecological risks associated with CC impacts on AS forests, with a particular focus on protective forests and their capacity to deliver Eco-DRR services under changing climatic conditions.

The database integrates **quantitative field and laboratory data, spatial datasets, and modelling tools** addressing multiple natural hazards and forest dynamics across spatial and temporal scales. These resources respond to key MOSAIC objectives, including: (i) collecting, harmonising, and sharing **Alpine-wide data** on historic and current climate-related hazards and forest disturbances; (ii) quantifying **past trends and future projections** under plausible climate scenarios; and (iii) **upgrading risk models** to account for CC-driven compound events affecting forest resilience and protective functions.

By making these resources openly accessible and interoperable, the deliverable supports knowledge transfer to professional actors, including **policymakers, risk managers, forest practitioners, and stakeholders** involved in forest living labs (FLLs). In doing so, it contributes to increasing awareness, fostering adaptive forest management approaches, and supporting evidence-based climate change mitigation and adaptation strategies within regional to transnational Alpine climate action plans.

1.3 Data Management, FAIR Principles, and Accessibility

All datasets and models included in this deliverable are documented following the project data management plan and the FAIR principles (Wilkinson et al., 2016), ensuring that project outputs are findable, accessible, interoperable, and reusable throughout and beyond the MOSAIC project lifetime. To this end, each resource is accompanied by standardized metadata describing its content, methods, spatial and temporal characteristics, access conditions, and recommended citation.

Alpine Space

MOSAIC

Adherence to the FAIR principles is implemented as follows:

- **Findable**

Each dataset and model is uniquely identified through a persistent identifier, preferably a **Digital Object Identifier (DOI)**, and described using rich, searchable metadata. These metadata enable discovery through the MOSAIC WebAtlas and external indexing services, facilitating visibility and traceability of project outputs.

- **Accessible**

Data and models are made openly accessible through stable repositories, using standard web protocols. The preferred solution is publication through dedicated services integrated within the MOSAIC **WebAtlas**, one of the main project results, which provides a centralized access point for project outputs. Where appropriate, links to established external open-access repositories are also supported to ensure continuity and broad dissemination.

- **Interoperable**

Resources are provided using open, **well-documented file formats and standardised metadata fields**, enabling integration with modelling platforms, GIS software, and other analytical tools. In particular, tables are provided preferably as csv, raster files as GeoTiff or ASCII, model codes as R or Python scripts or packages or through the upgrading of existing modeling platforms. The metadata framework is designed to support interoperability across datasets, models, and external platforms relevant to natural hazard and forest dynamics research.

- **Reusable**

Alpine Space

MOSAIC

Clear documentation, explicit licensing, and standardized citation guidelines are provided for each resource to enable correct interpretation and reuse. Metadata include methodological descriptions, assumptions, and known limitations, ensuring that datasets and models can be reliably reused in scientific, technical, and policy-oriented applications.

The preferred publication pathway for datasets and models is through an internal upload and curation within the MOSAIC WebAtlas (<https://alpineresilience.org/>), ensuring consistency, harmonization, and direct integration with project tools and visualization services. At the same time, the framework accommodates resources already published in established external repositories, which are referenced and linked within the MOSAIC database. This complementary approach ensures flexibility while maintaining a coherent and sustainable data infrastructure. Overall, this strategy guarantees long-term accessibility, citability, and traceability of MOSAIC outputs, supporting knowledge transfer, reproducibility, and reuse beyond the project duration.

1.4 Structure of the Deliverable

This deliverable is organised into two main parts. The first part provides an overview of the datasets, models, and derived products currently available in the MOSAIC database (*Chapter 2*). The second part introduces a standardized metadata submission framework designed to support consistent contributions from project partners and future extensions of the database (*Chapter 3*).

1.5 Related Documents

This deliverable is complemented by an Excel-based dataset inventory consisting of the table illustrated in *Chapter 2 Field and Lab Experiments Database*.

2. Field and Lab Experiments Database

2.1 Chapter introduction

This chapter presents the current inventory of datasets, models, and derived products included in the MOSAIC database under WP2, the core of **D.2.1.1**. The inventory provides an overview of resources generated within the project as well as selected external datasets and models that are relevant for analysing CC-driven compound events affecting AS protective forests.

The database is structured to ensure **clarity, traceability, and interoperability** across heterogeneous data types, supporting both integration into the NAZCA modelling platform and dissemination through the MOSAIC WebAtlas. Each record represents a distinct dataset or model and is documented using a standardised set of descriptive fields to facilitate discovery, reuse, and correct citation.

2.2 Structure of the database table

The database is presented as a **tabular inventory** in which **each row corresponds to a single data record** (dataset, model, or derived product), and **each column describes a specific attribute** of that record. The fields are defined as follows:

- **Code:** Unique identifier synthesizing the development source, main topic, and data type of the record.
- **Source:** Origin of the record, distinguishing between resources developed within MOSAIC and those sourced from external initiatives.
- **Main Topic:** The primary thematic focus of the record (e.g. wildfires, species suitability, rockfalls).

Alpine Space

MOSAIC

- **Data Type:** Classification of the resource as either data (e.g. raster files, tables) or model/code (e.g. R or Python scripts, software packages).
- **Title:** A concise and descriptive title summarizing the content of the record.
- **Description:** A short explanation of the record, including its purpose, methodological approach, and/or novelty.
- **PP(s):** Project Partner(s) responsible for data collection, development, and/or curation.
- **Link to Data:** Persistent link or DOI to the repository where the record is stored, either within the MOSAIC WebAtlas or in an external open-access repository.
- **Reference:** Bibliographic reference and/or DOI to publications describing or using the record, if available.

This standardized structure ensures consistency across contributions and supports both human interpretation and machine-readable integration with the MOSAIC web atlas.

Alpine Space

MOSAIC

2.3 The Database

Code	Source	Main Topic	Data Type	Title	Description	PP(s)	Link to Data	Reference
M_Win01_Rast	MOSAIC	Windthrows	Data (raster files)	<i>The protective effect of a windthrown forest: biological legacies and rockfalls</i>	The dataset uses the rockyfor3d model to assess the protective effect of a windthrown forest. Three scenarios are implemented: no forest, pre-storm forest conditions, and windthrown forest (i.e., post-storm conditions). The data are composed of three rasters, each representing one scenario. The rasters show the number of passages in the study area. The resolution is 2x2 meters. The dataset reports a new, innovative procedure for precisely assessing forest vulnerability to wind using remote-sensed data (LiDAR) and incorporating the ForestGALES model. This method automatically maps forest wind vulnerability at a high resolution (20x20 meters) and provides an example of a regional-scale application in the municipality of Rocca Pietore, Italy. The maps are rasters with a resolution of 20x20 meters.	PP4 (UNIPD)		<u>Costa, M., Marchi, N., Bettella, F., Bolzon, P., Berger, F., & Lingua, E. (2021). Biological Legacies and Rockfall: The Protective Effect of a Windthrown Forest. Forests, 12(9), 1141. https://doi.org/10.3390/f12091141</u>
M_Win02_Rast	MOSAIC	Windthrows	Data (raster files)	<i>Maps of forest wind vulnerability from remote sensed data through a novel semi-automatic methodology</i>	The dataset reports a new, innovative procedure for precisely assessing forest vulnerability to wind using remote-sensed data (LiDAR) and incorporating the ForestGALES model. This method automatically maps forest wind vulnerability at a high resolution (20x20 meters) and provides an example of a regional-scale application in the municipality of Rocca Pietore, Italy. The maps are rasters with a resolution of 20x20 meters.	PP4 (UNIPD)		<u>Baggio, T., Costa, M., Marchi, N., Locatelli, T., & Lingua, E. (2025). Improve the estimation of forest wind vulnerability through remote sensed data: a new methodology. Environmental Modelling & Software, 106825. https://doi.org/10.1016/j.envsoft.2025.106825</u>

Alpine Space

MOSAIC

Code	Source	Main Topic	Data Type	Title	Description	PP(s)	Link to Data	Reference
M_Win03_Mod	MOSAIC	Windthrows	Model	<i>A semi automatic methodology to assess forest wind vulnerability from remote sensed data</i>	The model is an innovative procedure that uses remote-sensed data (LiDAR) and incorporates the ForestGALES model to precisely assess forest vulnerability to wind. This method automatically maps forest wind vulnerability at a high resolution.	PP4 (UNIPD)	https://github.com/TommBagg/CWS_calculation_fgr	Baggio, T., Costa, M., Marchi, N., Locatelli, T., & Lingua, E. (2025). Improve the estimation of forest wind vulnerability through remote sensed data: a new methodology. Environmental Modelling & Software, 106825. https://doi.org/10.1016/j.envsoft.2025.106825
M_Fir01_Mod	MOSAIC	Wildfires	Model	<i>Models for post-fire applied nucleation under different restoration scenarios to improve degraded ecosystems - Bourra case study</i>	The dataset provides R code for developing a machine learning correlative model (Bayesian Additive Regression Tree, BART) at the landscape scale to model the occurrence of natural tree regeneration. Its goal is to provide a set of applied nucleation datasets for prioritizing post-fire intervention practices. Together with R code, the dataset comes with a table with data input for the Bourra site (Verrayes, Aosta, Italy).	PP3 (UNITO)		Mantero, G., Anselmetto, N., Morresi, D., Meloni, F., Bolzon, P., Lingua, E., ... & Marzano, R. (2024). Modeling post-fire regeneration patterns under different restoration scenarios to improve forest recovery in degraded ecosystems. Forest Ecology and Management, 551, 121520. https://doi.org/10.1016/j.foreco.2023.121520
M_Fir02_Rast	MOSAIC	Wildfires	Data (raster files)	<i>Applied nucleation maps of post-fire management under different restoration scenarios to improved</i>	The dataset provides raster maps for the simulations of the machine learning correlative model (Bayesian Additive Regression Tree, BART). They include a set of applied nucleation datasets for prioritizing post-fire intervention practices (EPSG:32632, resolution of 10m). Simulations for the Bourra site (Verrayes, Aosta, Italy).	PP3 (UNITO)	https://doi.org/10.6084/m9.figshare.23619018	Mantero, G., Anselmetto, N., Morresi, D., Meloni, F., Bolzon, P., Lingua, E., ... & Marzano, R. (2024). Modeling post-fire regeneration patterns under different restoration scenarios to improve forest recovery in degraded ecosystems. Forest Ecology and Management, 551, 121520. https://doi.org/10.1016/j.foreco.2023.121520

Alpine Space

MOSAIC

Code	Source	Main Topic	Data Type	Title	Description	PP(s)	Link to Data	Reference
				<i>degraded ecosystems - Bourra case study</i>				
M_Fir03_Rast	MOSAIC	Wildfires	Data (raster files)	<i>Modeling post-fire regeneration patterns under different restoration scenarios to improve forest recovery in degraded ecosystems - Mompantero case study</i>	The dataset provides raster maps for the simulations of the machine learning correlative model (Bayesian Additive Regression Tree, BART). They include a set of applied nucleation datasets for prioritizing post-fire intervention practices (EPSG:32632, resolution of 10m). Simulations for the Mompantero site (Turin, Italy).	PP3 (UNITO)		Mantero, G., Anselmetto, N., Morresi, D., Meloni, F., Bolzon, P., Lingua, E., ... & Marzano, R. (2024). Modeling post-fire regeneration patterns under different restoration scenarios to improve forest recovery in degraded ecosystems. <i>Forest Ecology and Management</i> , 551, 121520. https://doi.org/10.1016/j.foreco.2023.121520

Alpine Space

MOSAIC

Code	Source	Main Topic	Data Type	Title	Description	PP(s)	Link to Data	Reference
E_Suit01_Rast	External	Species Suitability	Data (raster files)	<i>Species distribution models built with local species data and regional climate models for the European Alps</i>	Collection of raster related to the historic and future distribution of 22 tree species for Piedmont Region, Italy and the European Alps extent produced by using correlative species distribution models. Rasters are projected to EPSG:3034 with a spatial resolution of 1km. Files are provided as both binary maps (0 = absence, 1 = presence) or continuous probability of presence (0-1).	PP3 (UNITO)	https://doi.org/10.6084/m9.figshare.26232551	Anselmetto, N., Morresi, D., Barbarino, S., Loglisci, N., Betts, M. G., & Garbarino, M. (2025). Species distribution models built with local species data perform better for current time, but suffer from niche truncation. <i>Agricultural and Forest Meteorology</i> , 362, 110361. https://doi.org/10.1016/j.agrformet.2024.110361
M_Suit02_Tab	MOSAIC	Species Suitability		Suitability Maps (IIASA)		PP7 (IIASA)		
M_Suit03_Rast	MOSAIC	Species Suitability	Data (raster files)	<i>Modeled suitability map for current forest stands under different climate change intensities</i>	Projection of suitability under different climate change scenarios for the species that currently compose forest stands. These maps are designed to identify areas where the expected effects of climate change on species composition shifts are highest. The maps represent the percentage of the forest composition of a current stand that is made up of species	PP2 (ONF)		

Alpine Space

MOSAIC

Code	Source	Main Topic	Data Type	Title	Description	PP(s)	Link to Data	Reference
M_Suit04_Rast	MOSAIC	Species Suitability		Incompatibility Maps (IIASA/ONF)	capable of surviving under future climate conditions (only species representing at least 20% of the current stand are included). Values range from 0% (no species in the current stand are expected to persist) to 100% (all species in the current stand are expected to persist). The dataset covers all French Alpine forests.	PP2 (ONF) & PP7 (IIASA)		
Rock01_Tab	External	Rockfalls		Rockfall database in Alpine Space	ROCKtheALPS	-		

Alpine Space

MOSAIC

Code	Source	Main Topic	Data Type	Title	Description	PP(s)	Link to Data	Reference
M_Dist01_Mod	MOSAIC		Model	<i>High-dimensional detection of Landscape Dynamics: a Landsat time series-based algorithm for forest disturbance mapping and beyond</i>	R package for HILANDYN (High-dimensional detection of Landscape Dynamics) algorithm, which exploits spatial and spectral information provided by Landsat time series to detect forest disturbance dynamics. HILANDYN is a novel and unsupervised procedure for changepoint detection in high-dimensional time series to segment inter-annual time series into linear trends. The algorithm embeds a noise filter to remove spurious changepoints caused by residual spectral noise in the time series. FLAM is a process-based fire model that uses parameterization algorithm originally developed to link a fire model with dynamic global vegetation models. The key features implemented in FLAM include fuel moisture computation based on the Fine Fuel Moisture Code (FFMC) of the Canadian Forest Fire Weather Index (FWI), a procedure to calibrate spatial fire suppression efficiency, and the ability to include additional variables in fire modeling and prediction.	PP3 (UNITO)	https://github.com/dnatorres/hilandyn	Morresi, D., Maeng, H., Marzano, R., Lingua, E., Motta, R., & Garbarino, M. (2024). High-dimensional detection of Landscape Dynamics: a Landsat time series-based algorithm for forest disturbance mapping and beyond. <i>GIScience & Remote Sensing</i>, 61(1). https://doi.org/10.1080/15481603.2024.2365001
E_Fir04_Mod	External		Model	<i>Wildfire climate impacts and adaptation model (FLAM)</i>	Wildfire climate impacts and adaptation model (FLAM)	PP7 (IIASA)	https://iiasa.ac.at/mo-dels-tools-data/flamm	Krasovskii, A., Khabarov, N., Pirker, J., Kraxner, F., Yowargana, P., Schepaschenko, D., & Obersteiner, M. (2018). Modeling Burned Areas in Indonesia: The FLAM Approach. <i>Forests</i>, 9(7), 437. https://doi.org/10.3390/f9070437

Alpine Space

MOSAIC

Code	Source	Main Topic	Data Type	Title	Description	PP(s)	Link to Data	Reference
E_Suit05_Mod	External	Species Suitability	Model	LandClim				
E_Ava01_Mod	External	Avalanche	Model	<i>Modified Version of the FlowPy Model with Automatic Energy Line Angle Selection</i>	This code is designed for large-scale identification of forests providing a protective effect against avalanches. It is based on the FlowPy model and integrates an INRAE-developed methodology enabling the automatic determination of the travel angle used to model avalanche propagation. Unlike the original implementation, which relies on a user-defined travel angle, the modified model computes, for each pixel, the dimensionless area beneath the topographic profile (i.e. the area under the profile normalized by the elevation difference relative to the release point). A statistical relationship is then applied to derive a travel angle adapted to local topographic conditions. The core model is implemented in Python and can be used as a standalone tool. It is also	PP1 (INRAE) & PP2 (ONF)		

Alpine Space

MOSAIC

Code	Source	Main Topic	Data Type	Title	Description	PP(s)	Link to Data	Reference
					embedded within an R interface that enables the processing of multiple spatial extents, including automatic detection of potential release areas and execution of the Python model.			
M_Fir05_Tab	MOSAIC	Wildfires	Data (table file)	<i>Post-fire natural regeneration across the Alps</i>	The data record represents a large dataset covering the countries of the Alpine Space to investigate short- and medium-term post-fire regeneration dynamics (from one to twenty years after disturbance). The dataset encompasses 14 fires distributed in Austria, France, Italy, Slovenia, and Switzerland. The database is presented as a table format, containing data derived from field surveys, historical and multitemporal data, and remote sensing analysis. The dataset presents information about the single regeneration tree and environmental variables about the	All PPs	https://alpineresilience.org/data	-

Alpine Space

MOSAIC

Code	Source	Main Topic	Data Type	Title	Description	PP(s)	Link to Data	Reference
					plot. For the single plant we report: species, height class, number of forest regeneration stems. For each plot we report: GPS coordinates of the center, altitude, fire severity, soil cover, and other climatic and topographical variables.			
M_RS01_Rast	MOSAIC	Remote Sensing	Data (raster files)	Orthophoto and structure-from-motion digital surface model (SfM-DSM) of UAV surveys for avalanche risk characterisation		PP4 (UNIPD) & PP6 (BFW)		-
M_Fir06_Tab	MOSAIC	Wildfires	Data (table file)	Soil water content and drought index from Cosmic Rays Neutron Sensing	The table file contains daily data on (i) volumetric water content (VWC) in the soil obtained using CRNS (Cosmic Rays Neutron Sensing) proximal probes - which quantify the absorption of neutrons produced by cosmic rays by water in the environment - and	PP5 (DPC/SP)		-

Alpine Space

MOSAIC

Code	Source	Main Topic	Data Type	Title	Description	PP(s)	Link to Data	Reference
M_Ava02_Vect	MOSAIC	Avalanche and Rockfall	Data (vector files)		dimensionless drought index (“Finapp Dryness Index”) derived from VWC through appropriate inversion and rescaling on a dimensionless scale.			
				<i>Decision-support map of forest protection functions against avalanche and rockfall processes</i>	Maps of avalanche and rockfall extents in French public forests, designed to support decision-making in forest planning. The maps are primarily produced through modelling of natural phenomena (rockfalls and avalanches), selecting all areas that intersect public forests and potentially affect infrastructure or human assets. The raw modelling results are then processed and harmonized to produce comprehensive maps that enable forest managers to quickly identify areas where forests can provide a protective function. The final product includes functional zoning of the phenomena (departure vs. transit areas), the importance of the protected assets (allowing prioritization), and their type.	PP1 (INRAE) & PP2 (ONF)		

Alpine Space

MOSAIC

Code	Source	Main Topic	Data Type	Title	Description	PP(s)	Link to Data	Reference
M_Fir07_Tab	MOSAIC	Wildfires	Data (table file)	<i>FWI Projections for the Triveneto Region</i>	The analysis estimates the future evolution of the Fire Weather Index (FWI) using maximum temperature and daily precipitation. The observed data and the calculated FWI were used to train a Random Forest model, which was then applied to data from four EURO-CORDEX models, bias-corrected and downscaled to the Triveneto region, for the RCP 4.5 and RCP 8.5 scenarios, for ten station points. A file is produced for each model station, containing FWI estimates for the period 1990-2100.	PP5 (DPC/SL)		https://clima.arpa.veneto.it/cline_approfondimento.pdf https://clima.arpa.veneto.it/info

3. Standardized Metadata Submission Framework

3.1 Introduction

This chapter describes the **standardised submission form** used to document and publish datasets, models, and derived products in the MOSAIC WebAtlas. The form defines a common set of metadata fields to ensure consistency, transparency, and interoperability across contributions from different PPs.

The submission form is designed to support the implementation of the **FAIR principles** by enabling clear identification, standardized documentation of methods, explicit access and licensing conditions, and persistent citation of project outputs, and is consistent with the metadata collection of common scientific repositories such as Dryad, Zenodo, Figshare, etc. It also ensures that contributed resources can be efficiently integrated into the MOSAIC WebAtlas, while **remaining reusable beyond the project lifetime**.

All partners contributing data or models to the MOSAIC database are required to complete this form. Mandatory fields are clearly indicated, while optional fields allow contributors to provide additional contextual information when relevant.

Alpine Space

MOSAIC

3.2 Submission Form Fields

3.2.1 Identification

Title * (text box)

Description: Mandatory. A clear and descriptive title that summarizes the dataset, model, or derived product. Avoid acronyms unless they are widely recognised. A length of approximately 120 characters is recommended.

Example: Post-fire natural regeneration across the Alps

Data Code / ID (auto-generated)

Description: No user input required. The identifier is assigned by the moderators and synthesizes the development source (M = MOSAIC, E = External), the main topic (e.g. wildfires, windthrows, species suitability), and the data type (e.g. model, raster, table).

Example: M_Fir01_Mod

Authors and Affiliations * (text box)

Description: Mandatory. Full names of authors (preferably including ORCID identifiers), institutional affiliations, and roles if relevant.

Example: Maria Rossi (ORCID: 0000-0002-...), University of North Pole, Department of Ecology;
Thomas Schmidt (ORCID: 0123-4567-...), University of South Pole, Department of Ecology

Contact Person(s) * (text box)

Description: Mandatory. Name and stable email address of the person responsible for the dataset or model. A project role may be added if useful.

Example: Maria Rossi – Data Curator – maria.rossi@uninp.np

Alpine Space

MOSAIC

3.2.2 Description

Type * (drop-down menu + optional text box)

Options: Dataset | Model/Code | Derived Product

Description: Mandatory. Select the primary resource type and, if relevant, specify a short subtype.

Examples: Dataset – Raster time series; Model – Python code; Derived product – Gridded indicator

Abstract * (text box)

Description: Mandatory. A plain-language summary describing the purpose, methods (1–2 lines), geographic and temporal coverage, main variables, key quality notes, and intended reuse. Internal jargon should be avoided. Recommended length: 1,000–2,000 characters.

Recommended Visualization (text box)

Description: Optional. Short indication (1–3 lines) of the most informative visualization for the web atlas (e.g. map, time series, histogram, dashboard). A small PNG/JPEG thumbnail may be provided if available.

Keywords (3–5) * (drop-down menu)

Description: Mandatory. Select 3–5 concise keywords from the predefined list. If necessary, select “other” and specify a new keyword.

Example: Regeneration; Wildfire; Recovery; Post-fire management

Version Number * (text box + changelog)

Description: Mandatory. Provide a semantic version number (MAJOR.MINOR, e.g. 1.0) and a brief changelog describing updates since the previous version.

Alpine Space

MOSAIC

Date of Last Upload (calendar field)
Description: Mandatory. Date of the most recent update in ISO 8601 format (YYYY-MM-DD).
Example: 2025-10-30

3.2.3 Spatial and temporal scales

(Displayed only if the resource is spatial or a spatially derived product)

Coordinate Reference System / Projection * (text box)
Description: Mandatory. Provide the CRS using an EPSG code.
Example: EPSG:4326

Geographic Extent * (four text boxes)
Description: Mandatory if applicable. Bounding box in WGS84 (EPSG:4326): West, East, South, North.
Example: W: 5.6; E: 16.0; S: 43.0; N: 48.5

Spatial Resolution (text box)
Description: Optional. Smallest spatial unit represented, including units.
Examples: 1-km grid; 10-m pixels; 1:25,000 scale; 6-m radius plots

Temporal Extent (text box)
Description: Optional. Start and end dates following ISO 8601. Use a single start date for ongoing datasets.
Examples: 2018/2025; 2019-06-01/2020-10-31; 2015-01-01/

Alpine Space

MOSAIC

Temporal Resolution (text box)
Description: Optional. Time interval between observations, if applicable.
Examples: Daily; Monthly; Annual; Single observation

3.2.4 Methods

Data Collection / Modelling Methods * (multiple text boxes)
Description: Mandatory. Provide a concise but complete description of how the dataset or model was produced, sufficient to ensure transparency, reproducibility, and correct reuse. Include the following elements as applicable:

- Purpose and context
- Data sources and preprocessing
- Sampling or experimental design
- Data processing and quality control
- Modelling approach (for models or derived products), including software, calibration, and validation
- Assumptions, limitations, and sources of uncertainty
- References to detailed methodological descriptions (e.g. publications, repositories)

Data Structure (text box)
Description: Optional. Description of file structure, variables, units, CRS (if spatial), table schemas, and folder organisation. A README file and, for tabular data, a data dictionary are strongly recommended.
Example: One CSV file with 10,000 rows and 15 variables; each row represents a 1 km² grid cell; coordinates in EPSG:4326.

Alpine Space

MOSAIC

File Format, Access, and Licensing * (drop-down menus + text box)

Description: Mandatory. Specify:

- File formats (prefer open standards such as CSV, GeoTIFF, GeoPackage, NetCDF/HDF5)
- Access link or repository URL
- Copyright holder and license (e.g. CC BY 4.0 for data; MIT/BSD/GPL for code)

Acceptance of Responsibilities (tick box)

Description: Mandatory. Confirmation that the contributor takes responsibility for the accuracy, ownership, and licensing of the submitted resource.

3.2.5 References

Related Publications (text box)

Description: Optional. Full bibliographic references (including DOIs where available) to publications describing or using the dataset or model.

DOI (automatically assigned)

Description: No user input required. A concept DOI will be assigned to the record, with version-specific DOIs for individual releases where applicable.

4. Conclusions and Outlook

This deliverable documents the first release of the MOSAIC database developed under WP2 NAZCA, providing a **harmonised and open-access collection** of datasets, models, and derived products addressing CC-driven compound events affecting AS protective forests. By integrating empirical data, spatial datasets, and modelling tools within a common framework, the deliverable establishes a critical foundation for the NAZCA modelling platform and for evidence-based climate adaptation and disaster risk reduction strategies in the AS.

Transnational added value. Climate-related hazards and their impacts on forest ecosystems do not stop at administrative or national borders. In the AS, shared climatic drivers, interconnected ecosystems, and transboundary socio-economic systems shaped by centuries of land use require coordinated approaches to risk assessment and adaptation planning of Eco-DRR. The MOSAIC database provides clear **transnational added value** by harmonising datasets and models originating from multiple countries (Italy, France, Austria, Switzerland, Slovenia, Germany), institutions (the different PPs), and disciplinary backgrounds into a **coherent and interoperable framework**. Through standardized metadata, persistent identifiers (i.e., DOI), and OA publication, the database enables **cross-border analyses** of hazard dynamics, forest vulnerability, and protective functions. This transnational perspective supports the development of shared knowledge bases, facilitates comparison of management strategies across regions, and strengthens cooperation among Alpine stakeholders, contributing to a common understanding of climate risks and responses.

Replicability in other regions. The approach adopted in this deliverable is designed to be replicable beyond the AS. The database structure, metadata framework, and submission procedures are generic and scalable, allowing their application to other mountain regions or

MOSAIC

landscapes exposed to CC-related hazards. By adhering to **FAIR** principles and relying on open standards and repositories, the MOSAIC database provides a **blueprint** for establishing similar platforms in different geographical and institutional contexts. This replicability enhances the broader relevance of MOSAIC, enabling the transfer of methods, tools, and data management practices to other regions facing comparable challenges related to CC, compound and cascade events, and Eco-DRR.

Key achievements of deliverable D.2.1.1. Deliverable D.2.1.1 achieves several key objectives. First, it establishes a **comprehensive inventory** of field and laboratory datasets, spatial products, and models relevant to CC-related compound events affecting AS protective forests. Second, it defines and implements a **standardised metadata framework** that ensures consistency, transparency, and interoperability across contributions. Third, it provides a **clear workflow** for data submission, documentation, and publication, facilitating ongoing contributions by project partners and integration into the MOSAIC WebAtlas. Together, these achievements lay the groundwork for sustained data sharing, model integration, and knowledge generation within WP2 NAZCA and across the MOSAIC project.

Contribution to scientific outputs. The MOSAIC database supports scientific research by providing **high-quality, well-documented datasets and models** that enable reproducible analyses of CC impacts, hazard dynamics, and forest ecosystem responses. By making these resources openly accessible and citable, the deliverable facilitates their use in peer-reviewed publications, comparative studies, and methodological developments. The integration of empirical data with modelling tools also fosters interdisciplinary research linking climate science, forest ecology, natural hazards, and socio-ecological systems.

Alpine Space

MOSAIC

Contribution to Policy Uptake. By translating complex data and models into accessible and interoperable resources, the database enhances the **uptake of scientific knowledge in policy and practice**. The integration of datasets into the MOSAIC WebAtlas directly supports decision-makers, risk managers, and forest practitioners in evaluating climate-related risks and adaptation options. This contributes to the development of evidence-based climate action plans, adaptive forest management strategies, and ecosystem-based disaster risk reduction policies at regional and transnational levels.

Contribution to long-term project legacy. Beyond the project duration, the MOSAIC database represents a lasting legacy by ensuring **long-term availability, traceability, and reuse of project outputs**. The use of persistent DOIs, OA repositories, and standardized metadata supports continued access and updates after project completion. By embedding the database within the MOSAIC WebAtlas and aligning it with existing Alpine initiatives, this deliverable contributes to sustaining a shared knowledge infrastructure that can support future research, policy development, and climate adaptation efforts.

5. References

Bastit, F., Brunette, M., & Montagné-Huck, C. (2023). Pests, wind and fire: A multi-hazard risk review for natural disturbances in forests. *Ecological Economics*, 205, 107702.

Cutter, S. L. (2018). Compound, cascading, or complex disasters: what's in a name?. *Environment: science and policy for sustainable development*, 60(6), 16-25.

Dorren, L. K., Berger, F., Imeson, A. C., Maier, B., & Rey, F. (2004). Integrity, stability and management of protection forests in the European Alps. *Forest ecology and management*, 195(1-2), 165-176.

IPCC, 2022. *Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press. Cambridge University Press, Cambridge, UK and New York, NY, USA, 3056 pp., doi:10.1017/9781009325844.

Moos, C., Stritih, A., Teich, M., & Bottero, A. (2023). Mountain protective forests under threat? an in-depth review of global change impacts on their protective effect against natural hazards. *Frontiers in Forests and Global Change*, 6, 1223934.

Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., ... & Mons, B. (2016). The FAIR Guiding Principles for scientific data management and stewardship. *Scientific data*, 3(1), 1-9.